

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Application of Robert A. Holton et al.
Serial No. 09/063,477
Filed April 20, 1998
For PROCESS FOR THE SELECTIVE DERIVATIZATION OF TAXANES
Examiner Ba K. Trinh

TO THE COMMISSIONER OF PATENTS AND TRADEMARKS

SIR:

DECLARATION OF DAVID J. PROCTER UNDER 37 C.F.R. 1.608(b)

I, David J. Procter, declare and state as follows:

1. At the time of the invention, I was a post doctorate student at Florida State University in Tallahassee, Florida, where I conducted research in Dr. Robert Holton's laboratory in the area of Synthetic Organic, Biorganic, and Organometallic Chemistry.

2. I am not an inventor of "Process For The Selective Derivatization Of Taxanes" of application no. 09/063,477.

3. During my tenure at Florida State University, I worked in the same laboratory as Zhuming Zhang. Zhuming Zhang worked in close proximity to me, and we routinely exchanged information about our experimental results. I observed Zhuming Zhang, Paul A. Clarke and Dr. Robert Holton reduce to practice the "Process For The Selective Derivatization Of Taxanes" before May 21, 1997.

4. Before May 21, 1997, I observed Zhuming Zhang conduct the "Attempt to protect C(7)OH by cbz" experiment, the "Attempt to protect C(10) H by (CH₃CO)-O" experiment, and the "Generation of baccatin III from 10DAB" experiment as documented on laboratory notebook pages 45, 49 and 67, respectively (Exhibits A-C). I recall that Zhuming Zhang immediately informed me of the results of these experiments so that I could use these processes to significantly simplify my own experimental research. Once Zhuming Zhang told me of his discoveries, I used these processes in preparing other taxane derivatives. The processes discovered by Zhuming Zhang eliminated about five process steps from my own experimental research in preparing the taxane derivatives.


5. After observing Zhuming Zhang conduct the "Attempt to protect C(10) H by (CH₃CO)-O" experiment shown on his laboratory notebook page 49, I used his process in my own research. For example, I conducted the "Selective Protection of 10-Hydroxyl - Formation of 10-Allyloxycarbonate" experiment shown on laboratory notebook pages 177 and 178 (Exhibit D). I added 0.137 ml of diallylpyrocarbonate to 30 mg of 10-deacetyl baccatin III in tetrahydrofuran solvent at room temperature and allowed the mixture to react while adding additional diallylpyrocarbonate at 21, 23, and 26 hours. The final reaction mixture revealed the formation of 10-allyloxycarbonate in about 63% of the reaction product.

6. After observing Zhuming Zhang conduct the "Generation of baccatin III from 10 DAB" experiment shown on his laboratory notebook page 67, I used his process in my own research. For example, I conducted the "Direct Acetylation of 10-DAB - Prepn. of B-III (ZnCl₂, 0°C)" experiment shown on laboratory notebook pages 167 and 168 shortly after his discovery (Exhibit E). I added 4 ml of acetic anhydride to 113 mg of 10-deacetyl baccatin III and 0.42 ml ZnCl₂ in tetrahydrofuran solvent at room temperature. Upon purification, 94.8 mg baccatin III was recovered, which was about 78% yield.


7. I further declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further, that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under 18 U.S.C. 1001, and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

1/8/00

Date


David J. Procter

Attempt to protect C(7) OH by Cbz



Materials used	Fw	AMT
(1) 10 DAB	544	3 mg (0.00551 mmol)
(2) dibenzyl dicarbonate (97%)	286.29	53 mg (38.9 μ , 0.984 mmol)
(3) THF		0.5 mL


Procedure: same as Page 44, except THF was used in stead of CH_2Cl_2 .

The rxn proceeded smoothly, TLC indicated the formation of product. The rxn was left for overnight.

Med:CHCl ₃ 1:9	Med:CHCl ₃ 1:9
0.5 h	overnight

This rxn almost go to completion,
very, very promising !!


¹H NMR @ 2-45-1 mixture of crude



So Cbz attached to C(10) OH

What a surprise ???!

high yield is obtained ($\geq 90\%$)



Materials used

FW

AMT

(1) 10DAB

544

3 mg (0.00551 mmol)

(2) Ac_2O

102.09

10 μL (20 eq., 0.11 mmol) $d = 1.082$

(3) THF



0.5 mL

Procedure: the rxn of 10DAB with $(\text{PhCH}_2\text{O})_2\text{C}=\text{O}$ suggested that similar reaction could take place between 10DAB + Ac_2O


To a solution of 10DAB in THF was added Ac_2O under Nitrogen. The reaction mixture was stirred at room temperature and monitored by TLC. TLC indicated the slow process of the rxn and a faint amount of product formed. At this stage it is not clear whether the rxn is proceeding or not. So 100 μL more Ac_2O was added. After 2h,

TLC

visible rxn was observed, a single spot shows up in high R_f value. After overnight, $\geq 80\%$ conversion was



1.5h after more Ac_2O was added



observed. At this stage, this rxn was stopped by evaporating the solvent away and taken by ^1H NMR

crude ^1H -NMR 2-49-1 indicated

$\geq 80\%$ - Baccatin III + 10DAB + small amount of (7-Ac²10DAB) (by evaporating the solvent?)

2-49-2 pure Baccatin III.

Procedure: To a THF solution of 10DAB + $ZnCl_2$ was added Al_2O_3 under N_2 . The solution was stirred at room temperature and monitored by TLC.

actually 18 mg $ZnCl_2$ (may have a little bit of H_2O)

EA: Hex 8:1

SM

0.5h

EA: Hex 8:1

SM

1h

EA: Hex 8:1

SM

1.5h

Hemochrom 1:7

SM

3h

2-67-2 crude mixture
Major baccatin III + small
amount 7 α H-diacetyl
+ oxetane ring opened
product.

Low temperature experiment is recommended.

137

DIRECT ACETYLATION OF 10-DAB - PREPN. OF B-III (ZnCl₂, 0°C).

TRANSFORMATION —

QUANTITIES


	QUANT:	MOL. WT:	MMOLS:	EQUIV:	SOURCE:
+ slight impurity +	10-DAB (recovered)	~113mg	544	0.21	1
d=1.052	Ac ₂ O	4ml	—	—	—
	ZnCl ₂ (0.5M THF)	0.42ml	—	0.21	1
	THF	2ml	—	—	—

METHOD — 1/2 Ac₂O (4ml) under N₂ was added ZnCl₂ in THF (0.42ml) to the resulting soln added to a soln. of 10-DAB (113mg) in THF (2ml) at 0°C.

Rxn. MONITORING —

On at 12.24 Fri.

50 mins.



80% EtAc/Hex

UV/Anisaldehyde

+ 1 mix of ZnCl₂ (0.42ml) added.

3h/ - trace of 10-DAB remaining.

4h/ - trace still remaining

rxn. mix. poured gradually into 45ml (NaHCO₃) added.

Aq. layer extracted with

EtAc (3x5ml)

dried (Na₂SO₄)

& concd. in vacuo => DJPI-167-

82.0

COMMENTS

Crude looks good

~10% bis acetate (+ trace of oxetane ring = opened product).

otherwise clean.

(60 mg, 11 %).

Purification
L

FLASH (50% EA/HEX)


[13-20] = 7 lbs. orotate

[22-45] = 7 BIII (94.8 mg, ~78 %)

→ DTPI-167-82-1

SELECTIVE PROTECTION OF 10-HYDROXYL- FORMATION OF 10-ALLYLOXYCARBONATE


{see 22 74}

TRANSFORMATIONQUANTITIES

	QUANT:	MOL. WT:	MMOLS:	EQUIV:	SOURCE:
10-DAB	30mg	544	0.055	1	/
d = 1.121 allyl pyrocarbonate	0.137ml	186.17	0.827	15	ALDRICH
THF	0.5ml	/	/	/	/

METHOD

1/2 a soln. of 10-DAB (30mg) in THF (0.5ml) at RT was added allyl pyrocarbonate (0.137ml, 15eq) & the rxn. mixture stirred at RT for h & the prod. followed by TLC.

REN. MONITORING

2h - rxn. proceeding slowly.

16h. rxn. gone ~50%

21h, 5eq. more of $(\sim 0.7)_2$ added.

23h, THF (0.5ml) & 10eq. reagent added (now identical conditions to ZZ).

26h, 10eq. more added (40eq. in total).

40h, rxn. still not complete.

W. Up rxn. mix. added to FLASH col. washed (50% EA/Hex).

RECOL \swarrow [2-5] \Rightarrow des. prod. + 10 DAB

FLASH (40% EA/Hex)

[4-7] \Rightarrow product (virtually pure).

(22 mg, ~63 %) \Rightarrow DJPI-177-87.0

[8-11] \Rightarrow 10-DAB.

(10.4 mg, ~32 %)

NEXT TIME

Recovered reagent DJPI-177-87.1

ALORICH reagent DJPI-177-87.2

- var. off. T₁

- rxn. mixture with 10% EA/Hex.

- rxn. mixture with 10% EA/Hex.